arXiv:1104.4099v1 [math.CO] 20 Apr 2011

Diagonalization of the Matrices of the Multinomial Descent and Multinomial Inversion Statistics on the Symmetric Group

Hery Randriamaro*
Fachbereich Mathematik und Informatik
Philipps-Universität Marburg
D-35032 Marburg
`herand@mathematik.uni-marburg.de`

January 20, 2013

Abstract

In the work of Varchenko, Zagier, Thibon, and Reiner, Saliola, Welker, linear algebraic properties of the multiplication map on the group algebra of the group algebra element are studied, which is the sum over all permutations weighted by $q^{\mathtt{inv}}$, $q^{\mathtt{maj}}$, `inv`. Here q is a variable, and `inv` and `maj` are the classical statistics inversion and major index. We define a multinomial descent statistic $\mathtt{des}_{\mathtt{X}}$ and a multinomial inversion statistic $\mathtt{inv}_{\mathtt{X}}$. These new defined statistics are the multinomial expressions of the classical statistics descent `des` and inversion. We determine the spectrum and the multiplicity of each element of the spectrum of the analogously defined multiplication map on the group algebra for both $\mathtt{des}_{\mathtt{X}}$ and $\mathtt{inv}_{\mathtt{X}}$. As corollaries we deduce the spectrum and the multiplicity of each element of the spectrum of the defined multiplication map on the group algebra for the statistics `des`, `maj` and `inv`.

1 Introduction

As usual we write $\mathcal{S}_n$ for the set of permutations of $[n] := \{1, 2, \dots, n\}$, $n \geq 1$. We consider for a permutation statistic $s : \mathcal{S}_n \to \mathbb{R}[X_1, \dots, X_k]$ the element $\mathsf{S} := \sum_{\sigma \in \mathcal{S}_n} s(\sigma)\sigma$ of the group algebra $\mathbb{R}[X_1, \dots, X_k][\mathcal{S}_n]$. The element S operates on $\mathbb{R}[X_1, \dots, X_k][\mathcal{S}_n]$ by left-multiplication as a linear mapping. The matrix of this linear mapping relativ to the standard basis $(\sigma)_{\sigma \in \mathcal{S}_n}$ is the matrix $\big(s(\sigma\tau^{-1})\big)_{\sigma, \tau \in \mathcal{S}_n}$.
We recall that for $\sigma \in \mathcal{S}_n$:

- its descents set is $\mathtt{DES}(\sigma) := \{k \in [n-1] \mid \sigma(k) > \sigma(k+1)\}$.
- its inversions set is $\mathtt{INV}(\sigma) := \{(i, j) \mid i < j,\, \sigma(i) > \sigma(j)\}$.

We explore the matrix for the statistics $\mathtt{des}_{\mathtt{X}} : \mathcal{S}_n \to \mathbb{R}[X_1, \dots, X_{n-1}]$:

$$\mathtt{des}_{\mathtt{X}}(\sigma) := \sum_{i \in \mathtt{DES}(\sigma)} X_i$$

and the statistics $\mathtt{inv}_{\mathtt{X}} : \mathcal{S}_n \to \mathbb{R}[X_{1,2}, \dots, X_{n-1,n}]$:

$$\mathtt{inv}_{\mathtt{X}}(\sigma) := \sum_{(i,j) \in \mathtt{INV}(\sigma)} X_{i,j}.$$

We remark that $\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\sigma)\sigma$ is an element of the descent algebra of $\mathbb{R}[X_1, \dots, X_{n-1}][\mathcal{S}_n]$.

Definition 1. *Let $n \geq 1$. We choose a fixed linear order of $\mathcal{S}_n$ and build the following matrices whose rows and columns are indexed with the elements of $\mathcal{S}_n$ with respect to this order:*

$$\mathfrak{D}_{\mathtt{n}} := \big(\mathtt{des}_{\mathtt{X}}(\pi\tau^{-1})\big)_{\pi, \tau \in \mathcal{S}_n},$$

$$\mathfrak{I}_{\mathtt{n}} := \big(\mathtt{inv}_{\mathtt{X}}(\pi\tau^{-1})\big)_{\pi, \tau \in \mathcal{S}_n}.$$

---

*This research was supported by DAAD.

For a square matrix A we write $Sp(\mathrm{A})$ for its spectrum and $V_{\mathrm{A}}(a)$ for the multiplicity of the eigenvalue $a \in Sp(\mathrm{A})$.
We want to determine $Sp(\mathfrak{D}_{\mathtt{n}})$ resp. $Sp(\mathfrak{I}_{\mathtt{n}})$ and the multiplicity of the elements of $Sp(\mathfrak{D}_{\mathtt{n}})$ resp. $Sp(\mathfrak{I}_{\mathtt{n}})$.
By linear algebraic calculating we get:

$(n = 1)$ $Sp(\mathfrak{D}_{\mathtt{1}}) = \{0\}$ and $V_{\mathfrak{D}_{\mathtt{1}}}(0) = 1$.

$(n = 2)$ $Sp(\mathfrak{D}_{\mathtt{2}}) = \{X_1, -X_1\}$ and $V_{\mathfrak{D}_{\mathtt{2}}}(X_1) = 1$, $V_{\mathfrak{D}_{\mathtt{2}}}(-X_1) = 1$.

For $n \geq 3$, we prove the following theorem:

Theorem 1. *Let $n \geq 3$. Then $\mathfrak{D}_{\mathtt{n}}$ is diagonalizable and:*

$$Sp(\mathfrak{D}_{\mathtt{n}}) = \{\frac{n!}{2}\sum_{k=1}^{n-1} X_k,\ -(n-2)!\sum_{k=1}^{n-1} X_k,\ 0\}$$

with:

- $V_{\mathfrak{D}_{\mathtt{n}}}\big(\frac{n!}{2}\sum_{k=1}^{n-1} X_k\big) = 1,$
- $V_{\mathfrak{D}_{\mathtt{n}}}\big(-(n-2)!\sum_{k=1}^{n-1} X_k\big) = \binom{n}{2},$
- $V_{\mathfrak{D}_{\mathtt{n}}}(0) = n! - \binom{n}{2} - 1.$

As a direct application of the Theorem 1, we have the two following corollaries. We recall that:

$$\mathtt{des}(\sigma) := \#\mathtt{DES}(\sigma) \text{ is the number of the descents of } \sigma,$$

$$\mathtt{maj}(\sigma) := \sum_{i \in \mathtt{DES}(\sigma)} i \text{ is the major index of } \sigma.$$

Corollary 1. *Let $n \geq 1$ and $\mathsf{D}_{\mathsf{n}} := \big(\mathtt{des}(\pi\tau^{-1})\big)_{\pi,\tau \in \mathcal{S}_n}$. Then D_{n} is diagonalizable and we have:*

$(n = 1)$ $Sp(\mathsf{D}_1) = \{0\}$ *and* $V_{\mathsf{D}_1}(0) = 1$.

$(n = 2)$ $Sp(\mathsf{D}_2) = \{1, -1\}$ *and* $V_{\mathsf{D}_2}(1) = 1$, $V_{\mathsf{D}_2}(-1) = 1$.

$(n \geq 3)$ $Sp(\mathsf{D}_{\mathsf{n}}) = \{\binom{n}{2}(n-1)!,\ 0,\ -(n-1)!\}$ *and*

- $V_{\mathsf{D}_{\mathsf{n}}}(\binom{n}{2}(n-1)!) = 1,$
- $V_{\mathsf{D}_{\mathsf{n}}}\big(-(n-1)!\big) = \binom{n}{2},$
- $V_{\mathsf{D}_{\mathsf{n}}}(0) = n! - \binom{n}{2} - 1.$

Proof. Set $X_i = 1$ in Theorem 1. ☐

The following corollary can be seen as a non-exponential version of a result by Thibon ([4], Theorem 56) who studied the matrix with entries $q^{\mathtt{maj}(\sigma\tau^{-1})}$ with deep results from the theory of noncommutative symmetric functions ([2] and [5]). It would be interesting to understand the deeper connection between the two results.

Corollary 2. *Let $n \geq 1$ and $\mathsf{M}_{\mathsf{n}} := \big(\mathtt{maj}(\pi\tau^{-1})\big)_{\pi,\tau \in \mathcal{S}_n}$. Then M_{n} diagonalizable and we have:*

$(n = 1)$ $Sp(\mathsf{M}_1) = \{0\}$ *and* $V_{\mathsf{M}_1}(0) = 1$.

$(n = 2)$ $Sp(\mathsf{M}_2) = \{1, -1\}$ *and* $V_{\mathsf{M}_2}(1) = 1$, $V_{\mathsf{M}_2}(-1) = 1$.

$(n \geq 3)$ $Sp(\mathsf{M}_{\mathsf{n}}) = \{\binom{n}{2}\frac{n!}{2},\ 0,\ -\frac{n!}{2}\}$ *and*

- $V_{\mathsf{M}_{\mathsf{n}}}(\binom{n}{2}\frac{n!}{2}) = 1,$
- $V_{\mathsf{M}_{\mathsf{n}}}(-\frac{n!}{2}) = \binom{n}{2},$
- $V_{\mathsf{M}_{\mathsf{n}}}(0) = n! - \binom{n}{2} - 1.$

Proof. Set $X_i = i$ in Theorem 1. ☐

By linear algebraic calculating we get:

$(n=1)$ $Sp(\mathfrak{I}_1) = \{0\}$ and $V_{\mathfrak{I}_1}(0) = 1$.

$(n=2)$ $Sp(\mathfrak{I}_2) = \{X_{1,2}, -X_{1,2}\}$ and $V_{\mathfrak{I}_2}(X_{1,2}) = 1$, $V_{\mathfrak{I}_2}(-X_{1,2}) = 1$.

$(n=3)$ $Sp(\mathfrak{I}_3) = \{3X_{1,2} + 3X_{1,3} + 3X_{2,3},\ -X_{1,2} - 2X_{1,3} - X_{2,3},\ -X_{1,2} + X_{1,3} - X_{2,3},\ 0\}$ and

- $V_{\mathfrak{I}_3}(3X_{1,2} + 3X_{1,3} + 3X_{2,3}) = 1$,
- $V_{\mathfrak{I}_3}(-X_{1,2} - 2X_{1,3} - X_{2,3}) = 2$,
- $V_{\mathfrak{I}_3}(-X_{1,2} + X_{1,3} - X_{2,3}) = 1$,
- $V_{\mathfrak{I}_3}(0) = 2$.

For $n \geq 4$, we prove the following theorem:

Theorem 2. *Let $n \geq 4$. Then $\mathfrak{I}_{\mathtt{n}}$ is diagonalizable and:*

$$Sp(\mathfrak{I}_{\mathtt{n}}) = \{\frac{n!}{2} \sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} X_{i,j},\ -(n-2)! \sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} (j-i)X_{i,j},$$
$$-(n-3)! \sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} \Big(n - 2(j-i)\Big)X_{i,j},\ 0\}$$

with

- $V_{\mathfrak{I}_{\mathtt{n}}}(\frac{n!}{2}\sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} X_{i,j}) = 1$,
- $V_{\mathfrak{I}_{\mathtt{n}}}\big(-(n-2)!\sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} (j-i)X_{i,j}\big) = n-1$,
- $V_{\mathfrak{I}_{\mathtt{n}}}\big(-(n-3)!\sum_{\{(i,j)\in[n]^2 \ \mid \ i<j\}} \Big(n - 2(j-i)\Big)X_{i,j}\big) = \binom{n-1}{2}$,
- $V_{\mathfrak{I}_{\mathtt{n}}}(0) = n! - \binom{n}{2} - n$.

We note that an exponential version of $\mathfrak{I}_{\mathtt{n}}$ has been studied by Varchenko [10]. He was able to give a beautiful formula for its determinant. This exponential version has attracted considerable interest in various areas of mathematics (e.g.[1], [11]). As a direct application of the Theorem 2, we have the following corollary. This result was obtained in a recent independent work of Renteln ([7], Section 4.8). The integrality assertion of this corollary is a very special case of a theorem of Reiner, Saliola, Welker ([6], Theorem 1.4) which also inspired our investigation. We recall that:

$$\mathtt{inv}(\sigma) := \#\mathtt{INV}(\sigma) \text{ is the number of inversions of } \sigma.$$

Corollary 3. *Let $n \geq 1$ and $\mathsf{I}_{\mathsf{n}} := \big(\mathtt{inv}(\pi\tau^{-1})\big)_{\pi,\tau\in\mathcal{S}_n}$. Then I_{n} is diagonalizable and we have:*

$(n=1)$ $Sp(\mathsf{I}_1) = \{0\}$ *and* $V_{\mathsf{I}_1}(0) = 1$.

$(n=2)$ $Sp(\mathsf{I}_2) = \{1, -1\}$ *and* $V_{\mathsf{I}_2}(1) = 1$, $V_{\mathsf{I}_2}(-1) = 1$.

$(n=3)$ $Sp(\mathfrak{I}_3) = \{9,\ -4,\ -1,\ 0\}$ *and*

- $V_{\mathfrak{I}_3}(9) = 1$,
- $V_{\mathfrak{I}_3}(-4) = 2$,
- $V_{\mathfrak{I}_3}(-1) = 1$,
- $V_{\mathfrak{I}_3}(0) = 2$.

$(n \geq 4)$

$$Sp(\mathsf{I}_{\mathsf{n}}) = \{\frac{n!}{2}\binom{n}{2},\ -\frac{(n+1)!}{6},\ -\frac{n!}{6},\ 0\}$$

and

- $V_{\mathsf{I}_{\mathsf{n}}}(\frac{n!}{2}\binom{n}{2}) = 1$,
- $V_{\mathsf{I}_{\mathsf{n}}}(\frac{(n+1)!}{6}) = n-1$,
- $V_{\mathsf{I}_{\mathsf{n}}}(\frac{n!}{6}) = \binom{n}{2}$,
- $V_{\mathsf{I}_{\mathsf{n}}}(0) = n! - \binom{n}{2} - n$.

Proof. Set $X_{i,j} = 1$ in Theorem 2. □

2 Multinomial Version of the Theorem of Perron-Frobenius

For a polynomial P and a monomial M in $\mathbb{R}[X_1, \dots, X_k]$, we write $[M]P$ for the coefficient of M in P. For a square matrix A we write $E_{\mathsf{A}}(a)$ for the eigenspace of the eigenvalue $a \in Sp(\mathsf{A})$ and for a vector v we write $< \mathsf{v} >$ for the subspace generated by v.

Proposition 2.1. *Let* $n \geq 2$ *and* $\mathfrak{P}_{\mathfrak{n}} = \big(P_{i,j}\big)_{i,j \in [n]}$ *be a matrix of polynomials* $P_{i,j} \in \mathbb{R}[X_1, \dots, X_k]$ *such that:*

(a) $P_{i,j} \neq 0$ *and* $[X_1^{i_1} \dots X_k^{i_k}]P_{i,j} \geq 0$*,*

(b) there is a polynomial $P_n \in \mathbb{R}[X_1, \dots, X_k]$ *such that, for any* $i', i'' \in [n]$*,*

$$\sum_{j=1}^{n} P_{i',j} = \sum_{j=1}^{n} P_{i'',j} = P_n.$$

Then $P_n \in Sp(\mathfrak{P}_{\mathfrak{n}})$ *and* $E_{\mathfrak{P}_{\mathfrak{n}}}(P_n) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} >$.

Proof. We prove the assertion by induction on n. With simple calculations, we get the result for $n = 2$. Now we assume that the assertion is proven for $n - 1 \geq 2$.
Let us consider the following system of n equations:

$$\begin{array}{ccccccccc}
P_{1,1}x_1 & + & P_{1,2}x_2 & + & \dots & + & P_{1,n-1}x_{n-1} & + & P_{1,n}x_n & = & P_n x_1 \\
P_{2,1}x_1 & + & P_{2,2}x_2 & + & \dots & + & P_{2,n-1}x_{n-1} & + & P_{2,n}x_n & = & P_n x_2 \\
\vdots & & \vdots & & & & \vdots & & \vdots & & \vdots \\
P_{n-1,1}x_1 & + & P_{n-1,2}x_2 & + & \dots & + & P_{n-1,n-1}x_{n-1} & + & P_{n-1,n}x_n & = & P_n x_{n-1} \\
P_{n,1}x_1 & + & P_{n,2}x_2 & + & \dots & + & P_{n,n-1}x_{n-1} & + & P_{n,n}x_n & = & P_n x_n
\end{array}$$

On the last row, we have $P_{n,1}x_1 + P_{n,2}x_2 + \dots + P_{n,n-1}x_{n-1} = (P_n - P_{n,n})x_n$. By multiplying the n-1 first rows with $(P_n - P_{n,n})$, and after by replacing $(P_n - P_{n,n})x_n$ with $P_{n,1}x_1 + P_{n,2}x_2 + \dots + P_{n,n-1}x_{n-1}$ in the n-1 first rows, we obtain the following system of n-1 equations:

$$\begin{array}{ccccccccc}
P_{1,1}(P_n - P_{n,n})x_1 & + & \dots & + & P_{1,n-1}(P_n - P_{n,n})x_{n-1} & + & P_{1,n}Q & = & P_{n-1}x_1 \\
P_{2,1}(P_n - P_{n,n})x_1 & + & \dots & + & P_{2,n-1}(P_n - P_{n,n})x_{n-1} & + & P_{2,n}Q & = & P_{n-1}x_2 \\
\vdots & & & & \vdots & & \vdots & & \vdots \\
P_{n-1,1}(P_n - P_{n,n})x_1 & + & \dots & + & P_{n-1,n-1}(P_n - P_{n,n})x_{n-1} & + & P_{n-1,n}Q & = & P_{n-1}x_{n-1}
\end{array}$$

where $P_{n-1} = P_n(P_n - P_{n,n})$ and $Q = P_{n,1}x_1 + P_{n,2}x_2 + \dots + P_{n,n-1}x_{n-1}$.
We have, for $i \in [n-1]$:

- $$\begin{aligned}
\sum_{j=1}^{n-1} [x_j]\big(P_{i,1}(P_n - P_{n,n})x_1 + \dots + P_{i,n-1}(P_n - P_{n,n})x_{n-1} + P_{i,n}Q\big) \\
= \sum_{j=1}^{n-1} P_{i,j}(P_n - P_{n,n}) + \sum_{j=1}^{n-1} P_{i,n}P_{n,j} \\
= (P_n - P_{i,n})(P_n - P_{n,n}) + P_{i,n}(P_n - P_{n,n}) \\
= P_{n-1}.
\end{aligned}$$

- Let $j \in [n-1]$ and set

  $$P'_{i,j} = [x_j]\big(P_{i,1}(P_n - P_{n,n})x_1 + \dots + P_{i\,n-1}(P_n - P_{n,n})x_{n-1} + P_{i,n}Q\big).$$

  By (a) and the definition of $\mathfrak{P}_{\mathfrak{n}}$, we have that $0 \neq P_{i,j}(P_n - P_{n,n})$ is a polynomial with non-negative coefficients only. Again by (a), it follows that:

  ▷ $P'_{i,j} = P_{i,j}(P_n - P_{n,n}) + P_{i,n}P_{n,j} \neq 0$,

▷ $(P'_{i,j}\,,\, X_1^{i_1}\dots X_k^{i_k}) \geq 0$, for $i_1,\dots,i_k \in \mathbb{N}$.

By induction, the solution of the system (2) is $x_1 = x_2 = \cdots = x_{n-1}$. By replacing $x_2, x_3, \dots, x_{n-1}$ by x_1 in $P_{n,1}x_1 + P_{n,2}x_2 + \cdots + P_{n,n-1}x_{n-1} = (P_n - P_{n,n})x_n$, we have then $x_1 = x_n$. Hence the solution of this system (1) is $x_1 = x_2 = \cdots = x_n$.

Therefore, $P_n \in Sp(\mathfrak{P}_{\mathfrak{n}})$ and $E_{\mathfrak{P}_{\mathfrak{n}}}(P_n) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} >$. □

Corollary 2.2. *Let* $n \geq 2$ *and* $\mathfrak{P}_{\mathfrak{n}} = \big(P_{i,j}\big)_{i,j\in[n]}$ *be a matrix of polynomials* $P_{i,j} \in \mathbb{R}[X_1,\dots,X_k]$ *such that:*

(a) $P_{i,j} \neq 0$ *and* $[X_1^{i_1}\dots X_k^{i_k}]P_{i,j} \geq 0$, *for* $i \neq j$,

(b) $P_{i,i} = 0$, *for* $i \in [n]$,

(c) there is a polynomial $P_n \in \mathbb{R}[X_1,\dots,X_k]$ *such that, for any* $i', i'' \in [n]$,

$$\sum_{j=1}^{n} P_{i',j} = \sum_{j=1}^{n} P_{i'',j} = P_n.$$

Then $P_n \in Sp(\mathfrak{P}_{\mathfrak{n}})$ *and* $E_{\mathfrak{P}_{\mathfrak{n}}}(P_n) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} >$.

Proof. The proof is obvious for $n = 2$.
Let us consider $n > 2$. We have the following system of n equations:

$$\begin{array}{ccccccccccc}
0 & + & P_{1,2}x_2 & + & \dots & + & P_{1,n-1}x_{n-1} & + & P_{1,n}x_n & = & P_nx_1 \\
P_{2,1}x_1 & + & 0 & + & \dots & + & P_{2,n-1}x_{n-1} & + & P_{2,n}x_n & = & P_nx_2 \\
\vdots & & \vdots & & & & \vdots & & \vdots & & \vdots \\
P_{n-1,1}x_1 & + & P_{n-1,2}x_2 & + & \dots & + & 0 & + & P_{n-1,n}x_n & = & P_nx_{n-1} \\
P_{n,1}x_1 & + & P_{n,2}x_2 & + & \dots & + & P_{n,n-1}x_{n-1} & + & 0 & = & P_nx_n
\end{array}$$

On the last row, we have $P_{n,1}x_1 + P_{n,2}x_2 + \cdots + P_{n,n-1}x_{n-1} = P_nx_n$. By multiplying the n-1 first rows with P_n, and after by replacing P_nx_n with $P_{n,1}x_1 + P_{n,2}x_2 + \cdots + P_{n,n-1}x_{n-1}$ in the n-1 first rows, we obtain the following system of n-1 equations:

$$\begin{array}{ccccccc}
P_{1,n}P_{n,1}x_1 & + & \dots & + & (P_nP_{1,n-1} + P_{1,n}P_{n,n-1})x_{n-1} & = & P_n^2x_1 \\
(P_nP_{2,1} + P_{2,n}P_{n,1})x_1 & + & \dots & + & (P_nP_{2,n-1} + P_{2,n}P_{n,n-1})x_{n-1} & = & P_n^2x_2 \\
\vdots & & & & \vdots & & \vdots \\
(P_nP_{n-1,1} + P_{n-1,n}P_{n,1})x_1 & + & \dots & + & P_{n-1,n}P_{n,n-1}x_{n-1} & = & P_n^2x_{n-1}
\end{array}$$

where

$$\sum_{j=1}^{n-1}[x_j]\big(\sum_{k=1}^{n-1}[x_k](P_nP_{i,k} + P_{i,n}P_{n,k})\big) = P_n\sum_{k=1}^{n-1}P_{i,k} + P_{i,n}\sum_{k=1}^{n-1}P_{n,k} = P_n\sum_{k=1}^{n-1}P_{i,k} + P_nP_{i,n} = P_n^2.$$

Using Proposition 2.1 on the obtained system, we get $x_1 = x_2 = \cdots = x_{n-1}$ and then the desided result. □

Proposition 2.1 can also be applied to get more general results. As example, for a matrix of polynomials $\mathfrak{P}_{\mathfrak{n}} = \big(P_{i,j}\big)_{i,j\in[n]}$ such that:

(a) $[X_1^{i_1}\dots X_k^{i_k}]P_{i,j} \geq 0$,

(b) $P_{i,n} \neq 0$ and $P_{n,i} \neq 0$, for $i \neq n$,

(c) there is a polynomial $P_n \in \mathbb{R}[X_1,\dots,X_k]$ such that, for any $i', i'' \in [n]$,

$$\sum_{j=1}^{n} P_{i',j} = \sum_{j=1}^{n} P_{i'',j} = P_n.$$

Then $P_n \in Sp(\mathfrak{P}_{\mathfrak{n}})$ and $E_{\mathfrak{P}_{\mathfrak{n}}}(P_n) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} >$.

Now we apply Corollary 2.2 to the statistics $\mathtt{des}_{\mathtt{X}}$ and $\mathtt{inv}_{\mathtt{X}}$.

Lemma 2.3. *For $n \geq 3$, we have*

$$\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\sigma) = \frac{n!}{2} \sum_{k=1}^{n-1} X_k.$$

Proof. For $k \in [n-1]$, $[X_k]\big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\sigma)\big) = \#\{\sigma \in \mathcal{S}_n \,|\, k \in \mathtt{DES}(\sigma)\}$. It is clear that the following mapping is bijective:

$$\gamma : \left\{ \begin{array}{ccc} \{\sigma \in \mathcal{S}_n \,|\, k \in \mathtt{DES}(\sigma)\} & \to & \{\sigma \in \mathcal{S}_n \,|\, k \notin \mathtt{DES}(\sigma)\} \\ \tau = \begin{pmatrix} \dots & k & k+1 & \dots \\ \dots & \tau(k) & \tau(k+1) & \dots \end{pmatrix} & \mapsto & \gamma(\tau) = \begin{pmatrix} \dots & k & k+1 & \dots \\ \dots & \tau(k+1) & \tau(k) & \dots \end{pmatrix} \end{array} \right. ,$$

that means $\#\{\sigma \in \mathcal{S}_n \,|\, k \in \mathtt{DES}(\sigma)\} = \#\{\sigma \in \mathcal{S}_n \,|\, k \notin \mathtt{DES}(\sigma)\}$.
Since $\#\{\sigma \in \mathcal{S}_n \,|\, k \in \mathtt{DES}(\sigma)\} + \#\{\sigma \in \mathcal{S}_n \,|\, k \notin \mathtt{DES}(\sigma)\} = \#\mathcal{S}_n$, then $[X_j]\big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\sigma)\big) = \frac{n!}{2}$, and $\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\sigma) = \frac{n!}{2} \sum_{k=1}^{n-1} X_k$. ☐

From Corollary 2.2 and Lemma 2.3, we deduce that

$$\frac{n!}{2} \sum_{k=1}^{n-1} X_k \in Sp(\mathfrak{D}_{\mathfrak{n}}) \text{ and } E_{\mathfrak{D}_{\mathfrak{n}}}\Big(\frac{n!}{2} \sum_{k=1}^{n-1} X_k\Big) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} > . \tag{1}$$

Besides, from Lemma 2.3, we also directly get

$$\sum_{\sigma \in \mathcal{S}_n} \mathtt{des}(\sigma) = \frac{(n-1)n!}{2} \text{ and } \sum_{\sigma \in \mathcal{S}_n} \mathtt{maj}(\sigma) = \frac{n!}{2} \binom{n}{2}$$

that are differently calculated in other books ([9], Example 2.2.5 resp. Corollary 4.5.9 for example).

Lemma 2.4. *For $n \geq 4$, we have*

$$\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma) = \frac{n!}{2} \sum_{\{(i,j) \in [n]^2 \;|\; i<j\}} X_{i,j}.$$

Proof. For $i, j \in [n]$, $i < j$, $[X_{i,j}]\big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\big) = \#\{\sigma \in \mathcal{S}_n \,|\, (i,j) \in \mathtt{INV}(\sigma)\}$. It is clear that the following mapping is bijective:

$$\varphi : \left\{ \begin{array}{ccc} \{\sigma \in \mathcal{S}_n \,|\, (i,j) \in \mathtt{INV}(\sigma)\} & \to & \{\sigma \in \mathcal{S}_n \,|\, (i,j) \notin \mathtt{INV}(\sigma)\} \\ \tau = \begin{pmatrix} \dots & i & \dots & j & \dots \\ \dots & \tau(i) & \dots & \tau(j) & \dots \end{pmatrix} & \mapsto & \varphi(\tau) = \begin{pmatrix} \dots & i & \dots & j & \dots \\ \dots & \tau(j) & \dots & \tau(i) & \dots \end{pmatrix} \end{array} \right. ,$$

that means $\#\{\sigma \in \mathcal{S}_n \,|\, (i,j) \in \mathtt{INV}(\sigma)\} = \#\{\sigma \in \mathcal{S}_n \,|\, (i,j) \notin \mathtt{INV}(\sigma)\}$.
Since $\#\{\sigma \in \mathcal{S}_n \,|\, (i,j) \in \mathtt{INV}(\sigma)\} + \#\{\sigma \in \mathcal{S}_n \,|\, (i,j) \notin \mathtt{INV}(\sigma)\} = \#\mathcal{S}_n$, then $[X_{i,j}]\big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\big) = \frac{n!}{2}$, and $\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma) = \frac{n!}{2} \sum_{\{(i,j) \in [n]^2 \;|\; i<j\}} X_{i,j}$. ☐

From Corollary 2.2 and Lemma 2.4, we deduce that

$$\frac{n!}{2} \sum_{\{(i,j) \in [n]^2 \;|\; i<j\}} X_{i,j} \in Sp(\mathfrak{I}_{\mathfrak{n}}) \text{ and } E_{\mathfrak{I}_{\mathfrak{n}}}\Big(\frac{n!}{2} \sum_{\{(i,j) \in [n]^2 \;|\; i<j\}} X_{i,j}\Big) =< \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} > . \tag{2}$$

Besides, from Lemma 2.4, we also directly get

$$\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}(\sigma) = \frac{n!}{2} \binom{n}{2}$$

that is differently calculated in other books ([9], Corollary 1.3.10 for example).

3 Minimal Polynomial of the Multinomial Descent Statistic

To determine the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$ we need the both following lemmas:
Let $n \geq 3$ and $i, j \in [n]$. We write

$$\mathcal{S}_n^{i,j} := \{\sigma \in \mathcal{S}_n \mid \sigma^{-1}(j) - \sigma^{-1}(i) = 1\},$$
$$\mathcal{S}_n^{i\,-\,j} := \{\sigma \in \mathcal{S}_n \mid \sigma^{-1}(j) - \sigma^{-1}(i) > 1\}.$$

Let $n \geq 3$. We write

$$\mathfrak{d}_\mathfrak{n} = \sum_{k=1}^{n-1} X_k.$$

Lemma 3.1. *Let $n \geq 3$ and $i, j \in [n]$. Then*

- *if $j > i + 1$:*
$$\sum_{\sigma \in \mathcal{S}_n^{i,j}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{j,i}} X_{\sigma^{-1}(j)} \mathtt{des}_\mathtt{X}(\sigma^{-1}).$$
- *if $j = i + 1$:*
$$\sum_{\sigma \in \mathcal{S}_n^{i,(i+1)}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{(i+1),i}} X_{\sigma^{-1}(i+1)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) \ - X_i (n-2)! \mathfrak{d}_\mathfrak{n}.$$

Proof. Let us consider the following bijective mapping:

$$\kappa_{i,j} : \begin{cases} \mathcal{S}_n & \to & \mathcal{S}_n \\ \sigma = \begin{pmatrix} \dots & i & \dots & j & \dots \\ \dots & \sigma(i) & \dots & \sigma(j) & \dots \end{pmatrix} & \mapsto & \kappa_{i,j}(\sigma) = \begin{pmatrix} \dots & i & \dots & j & \dots \\ \dots & \sigma(j) & \dots & \sigma(i) & \dots \end{pmatrix} \end{cases}.$$

- Let $\sigma \in \mathcal{S}_n^{i\,j}$. Then we have the following simple facts:
  - ▷ $\sigma^{-1}(j) = \sigma^{-1}(i) + 1$,
  - ▷ $\sigma^{-1} \in \mathcal{S}_n^{\sigma^{-1}(i)\,-\,\sigma^{-1}(j)}$,
  - ▷ $\kappa_{i,j}(\sigma^{-1}) \in \mathcal{S}_n^{\sigma^{-1}(j)\,-\,\sigma^{-1}(i)}$,
  - ▷ $\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \mathtt{des}_\mathtt{X}\big(\kappa_{i,j}(\sigma^{-1})\big)$,
  - ▷ $\big(\kappa_{i,j}(\sigma^{-1})\big)^{-1} \in \mathcal{S}_n^{j,i}$.

  Thus
$$\sum_{\sigma \in \mathcal{S}_n^{i,j}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{i,j}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}\big(\kappa_{i,j}(\sigma^{-1})\big) = \sum_{\sigma \in \mathcal{S}_n^{j,i}} X_{\sigma^{-1}(j)} \mathtt{des}_\mathtt{X}(\sigma^{-1}).$$

- Let $\sigma \in \mathcal{S}_n^{i\,(i+1)}$. Again the following facts hold:
  - ▷ $\sigma^{-1} \in \mathcal{S}_n^{\sigma^{-1}(i),(\sigma^{-1}(i)+1)}$,
  - ▷ $\kappa_{i,i+1}(\sigma^{-1}) \in \mathcal{S}_n^{(\sigma^{-1}(i)+1),\sigma^{-1}(i)}$,
  - ▷ $\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \mathtt{des}_\mathtt{X}\big(\kappa_{i,i+1}(\sigma^{-1})\big) - X_i$,
  - ▷ $\big(\kappa_{i,i+1}(\sigma^{-1})\big)^{-1} \in \mathcal{S}_n^{(i+1),i}$.

  Thus
$$\sum_{\sigma \in \mathcal{S}_n^{i\,(i+1)}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{i,(i+1)}} X_{\sigma^{-1}(i)} \Big(\mathtt{des}_\mathtt{X}\big(\kappa_{i,i+1}(\sigma^{-1})\big) - X_i\Big)$$
$$= \sum_{\sigma \in \mathcal{S}_n^{i\,(i+1)}} X_{\sigma^{-1}(i)} \mathtt{des}_\mathtt{X}\big(\kappa_{i,i+1}(\sigma^{-1})\big) - \sum_{\sigma \in \mathcal{S}_n^{i,(i+1)}} X_{\sigma^{-1}(i)} X_i = \sum_{\sigma \in \mathcal{S}_n^{(i+1),i}} X_{\sigma^{-1}(i+1)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) - X_i \sum_{k=1}^{n-1} \sum_{\substack{\sigma \in \mathcal{S}_n^{i,(i+1)} \\ \sigma(k)=i}} X_k$$
$$= \sum_{\sigma \in \mathcal{S}_n^{(i+1),i}} X_{\sigma^{-1}(i+1)} \mathtt{des}_\mathtt{X}(\sigma^{-1}) - X_i (n-2)! \mathfrak{d}_\mathfrak{n}.$$

$\square$

Let $n \geq 3$. We write ι for the identity permutation of $\mathcal{S}_n$ and define:

$$\begin{array}{rccc} \Pi_n : & \mathcal{S}_n & \to & \mathbb{R}[X_1, \dots, X_{n-1}] \\ & \tau & \mapsto & \sum_{\sigma \in \mathcal{S}_n} \mathtt{des}_{\mathtt{X}}(\tau\sigma)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) \end{array}$$

Lemma 3.2. *Let $n \geq 3$ and $\tau \in \mathcal{S}_n$. Then*

$$\Pi_n(\tau) = \Pi_n(\iota) - \mathtt{des}_{\mathtt{X}}(\tau)(n-2)!\mathfrak{d}_{\mathfrak{n}}.$$

Proof. Every permutation $\tau \in \mathcal{S}_n$ can be written as $\tau = \tau_k \dots \tau_2\tau_1\iota$ for some $\tau_i \in \{(j\ j+1) \mid 1 \leq j \leq n-1\}$. We consider the following mapping:

$$\phi : \left\{ \begin{array}{ccc} \mathcal{S}_n & \to & \mathbb{R}[X_1, \dots, X_{n-1}] \\ \tau = \tau_k \dots \tau_2\tau_1\iota & \mapsto & \Big(\mathtt{des}_{\mathtt{X}}(\tau) - \mathtt{des}_{\mathtt{X}}(\tau_{k-1} \dots \tau_2\tau_1\iota)\Big) + \phi\Big(\tau_{k-1} \dots \tau_2\tau_1\iota\Big) \end{array} \right. ,$$

$$\text{where } \phi(\iota) = 0 \text{ and } \phi\Big((j\ j+1)\Big) = X_j.$$

We are going to prove that $\Pi_n(\tau) = \Pi_n(\iota) - \phi(\tau)(n-2)!\mathfrak{d}_{\mathfrak{n}}$.
The case $k = 0$ and $\tau = \iota$ is trivial.
For $k = 1$ and $\tau = (j\ j+1)$ we have:

$$\Pi_n\big((j\ j+1)\big) = \sum_{\sigma \in \mathcal{S}_n^{j\,-\,(j+1)}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) + \sum_{\sigma \in \mathcal{S}_n^{(j+1)\,-\,j}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) +$$

$$\sum_{\sigma \in \mathcal{S}_n^{j,(j+1)}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) + \sum_{\sigma \in \mathcal{S}_n^{(j+1),j}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) =$$

$$\sum_{\sigma \in \mathcal{S}_n^{j\,-\,(j+1)}} \mathtt{des}_{\mathtt{X}}(\sigma)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) + \sum_{\sigma \in \mathcal{S}_n^{(j+1)\,-\,j}} \mathtt{des}_{\mathtt{X}}(\sigma)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) +$$

$$\sum_{\sigma \in \mathcal{S}_n^{j,(j+1)}} \Big(\mathtt{des}_{\mathtt{X}}(\sigma) + X_{\sigma^{-1}(j)}\Big)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) + \sum_{\sigma \in \mathcal{S}_n^{(j+1),j}} \Big(\mathtt{des}_{\mathtt{X}}(\sigma) - X_{\sigma^{-1}(j+1)}\Big)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) =$$

$$\Pi_n(\iota) + \sum_{\sigma \in \mathcal{S}_n^{j,(j+1)}} X_{\sigma^{-1}(j)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) - \sum_{\sigma \in \mathcal{S}_n^{(j+1),j}} X_{\sigma^{-1}(j+1)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) =$$

$$\Pi_n(\iota) + \sum_{\sigma \in \mathcal{S}_n^{(j+1),j}} X_{\sigma^{-1}(j+1)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) - X_j(n-2)!\mathfrak{d}_{\mathfrak{n}} - \sum_{\sigma \in \mathcal{S}_n^{(j+1),j}} X_{\sigma^{-1}(j+1)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) =$$

$$\Pi_n(\iota) - \phi\Big((j\ j+1)\Big)(n-2)!\mathfrak{d}_{\mathfrak{n}}.$$

Now we assume that the assertion is proven for $k - 1 \geq 1$.
Let $k \geq 2$ and $\tau = \tau_k \dots \tau_2\tau_1\iota = (j\ j+1)\tau'$, where $\tau_k = (j\ j+1)$ and $\tau' = \tau_{k-1} \dots \tau_2\tau_1\iota$:

$$\Pi_n\big((j\,j{+}1)\tau'\big) = \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{j\,-\,(j+1)}\}} \mathtt{des}_{\mathtt{X}}\Big((j\,j{+}1)\tau'\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) + \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{(j+1)\,-\,j}\}} \mathtt{des}_{\mathtt{X}}\Big((j\,j{+}1)\tau'\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) +$$

$$\sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{j,(j+1)}\}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\tau'\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) + \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{(j+1),j}\}} \mathtt{des}_{\mathtt{X}}\Big((j\ j+1)\tau'\sigma\Big)\mathtt{des}_{\mathtt{X}}\Big(\sigma^{-1}\Big) =$$

$$\sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{j\,-\,(j+1)}\}} \mathtt{des}_{\mathtt{X}}(\tau'\sigma)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) + \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{(j+1)\,-\,j}\}} \mathtt{des}_{\mathtt{X}}(\tau'\sigma)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) +$$

$$\sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{j,(j+1)}\}} \Big(\mathtt{des}_{\mathtt{X}}(\tau'\sigma) + X_{\sigma^{-1}\tau'^{-1}(j)}\Big)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) + \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{(j+1),j}\}} \Big(\mathtt{des}_{\mathtt{X}}(\tau'\sigma) - X_{\sigma^{-1}\tau'^{-1}(j+1)}\Big)\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) =$$

$$\Pi_n(\tau') + \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{j,(j+1)}\}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) - \sum_{\{\sigma \in \mathcal{S}_n \,\mid\, \tau'\sigma \in \mathcal{S}_n^{(j+1),j}\}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) =$$

$$\Pi_n(\tau') + \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j),\tau'^{-1}(j+1)}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}) - \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j+1),\tau'^{-1}(j)}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_{\mathtt{X}}(\sigma^{-1}).$$

- If $\tau' \in \mathcal{S}_n^{j\,-\,(j+1)}$ i.e. $\tau'^{-1}(j+1) - \tau'^{-1}(j) > 1$ and $\mathtt{des}_\mathtt{X}\Big((j\ j+1)\tau'\Big) = \mathtt{des}_\mathtt{X}(\tau')$, then

  $$\sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j),\tau'^{-1}(j+1)}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j+1),\tau'^{-1}(j)}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_\mathtt{X}(\sigma^{-1})$$

  and

  $$\Pi_n\big((j\ j+1)\tau'\big) = \Pi_n(\tau') = \Pi_n(\iota) - \phi\Big((j\ j+1)\tau'\Big)(n-2)!\mathfrak{d}_\mathfrak{n}.$$

- If $\tau' \in \mathcal{S}_n^{(j+1)\,-\,j}$ i.e. $\tau'^{-1}(j) - \tau'^{-1}(j+1) > 1$ and $\mathtt{des}_\mathtt{X}\Big((j\ j+1)\tau'\Big) = \mathtt{des}_\mathtt{X}(\tau')$, then

  $$\sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j),\tau'^{-1}(j+1)}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j+1),\tau'^{-1}(j)}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_\mathtt{X}(\sigma^{-1})$$

  and, once again, we have

  $$\Pi_n\big((j\ j+1)\tau'\big) = \Pi_n(\tau') = \Pi_n(\iota) - \phi\Big((j\ j+1)\tau'\Big)(n-2)!\mathfrak{d}_\mathfrak{n}.$$

- If $\tau' \in \mathcal{S}_n^{j,(j+1)}$ i.e. $\tau'^{-1}(j+1) - \tau'^{-1}(j) = 1$ and $\mathtt{des}_\mathtt{X}\Big((j\ j+1)\tau'\Big) = \mathtt{des}_\mathtt{X}(\tau') + X_{\tau'^{-1}(j)}$, then

  $$\sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j),\tau'^{-1}(j+1)}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j+1),\tau'^{-1}(j)}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) \ - X_{\tau'^{-1}(j)}(n-2)!\mathfrak{d}_\mathfrak{n}$$

  and

  $$\Pi_n\big((j\ j+1)\tau'\big) = \Pi_n\big(\tau'\big) - X_{\tau'^{-1}(j)}(n-2)!\mathfrak{d}_\mathfrak{n} = \Pi_n(\iota) - \phi\Big((j\ j+1)\tau'\Big)(n-2)!\mathfrak{d}_\mathfrak{n}.$$

- If $\tau' \in \mathcal{S}_n^{(j+1),j}$ i.e. $\tau'^{-1}(j) - \tau'^{-1}(j+1) = 1$ and $\mathtt{des}_\mathtt{X}\Big((j\ j+1)\tau'\Big) = \mathtt{des}_\mathtt{X}(\tau') - X_{\tau'^{-1}(j+1)}$, then

  $$\sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j),\tau'^{-1}(j+1)}} X_{\sigma^{-1}\tau'^{-1}(j)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) = \sum_{\sigma \in \mathcal{S}_n^{\tau'^{-1}(j+1),\tau'^{-1}(j)}} X_{\sigma^{-1}\tau'^{-1}(j+1)}\mathtt{des}_\mathtt{X}(\sigma^{-1}) + X_{\tau'^{-1}(j+1)}(n-2)!\mathfrak{d}_\mathfrak{n}$$

  and

  $$\Pi_n\big((j\ j+1)\tau'\big) = \Pi_n\big(\tau'\big) + X_{\tau'^{-1}(j)}(n-2)!\mathfrak{d}_\mathfrak{n} = \Pi_n(\iota) - \phi\Big((j\ j+1)\tau'\Big)(n-2)!\mathfrak{d}_\mathfrak{n}.$$

Finally, we have

$$\begin{array}{rcl}
\Pi_n(\tau) & = & \Pi_n(\iota) - \phi(\tau)(n-2)!\mathfrak{d}_\mathfrak{n} \\
 & = & \Pi_n(\iota) - \Big(\Big(\mathtt{des}_\mathtt{X}(\tau) - \mathtt{des}_\mathtt{X}(\tau_{k-1}\ldots\tau_2\tau_1\iota)\Big) + \phi\Big(\tau_{k-1}\ldots\tau_2\tau_1\iota\Big)\Big)(n-2)!\mathfrak{d}_\mathfrak{n} \\
 & = & \Pi_n(\iota) - \Big(\mathtt{des}_\mathtt{X}(\tau) - \mathtt{des}_\mathtt{X}(\iota)\Big)(n-2)!\mathfrak{d}_\mathfrak{n} \\
 & = & \Pi_n(\iota) - \mathtt{des}_\mathtt{X}(\tau)(n-2)!\mathfrak{d}_\mathfrak{n}.
\end{array}$$

□

Now we are in position to determine the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$.
We write $\mathrm{I}_{n!}$ for the matrix identity of dimension $n!$.

Proposition 3.3. *Let $n \geq 3$. The minimal polynomial of $\mathfrak{D}_\mathfrak{n}$ is:*

$$X\big(X - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\big)\big(X + (n-2)!\mathfrak{d}_\mathfrak{n}\big).$$

Proof. Using Lemma 2.3 and Lemma 3.2, we get:

$$\Big(\mathfrak{D}_\mathfrak{n} - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\Big)\mathfrak{D}_\mathfrak{n}\Big(\mathfrak{D}_\mathfrak{n} + (n-2)!\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\Big) = \Big(\mathfrak{D}_\mathfrak{n} - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\Big)\Big(\Pi_n(\iota)\Big)_{\pi,\tau\in\mathcal{S}_n} = \Big(0\Big)_{\pi,\tau\in\mathcal{S}_n}.$$

Hence the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$ divides $X\big(X - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\big)\big(X + (n-2)!\mathfrak{d}_\mathfrak{n}\big)$.
Since $\Big((\mathfrak{D}_\mathfrak{n} - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!})\mathfrak{D}_\mathfrak{n}\Big)_{\pi,\pi} = \Pi_n$ resp. $\Big(\mathfrak{D}_\mathfrak{n}\big(\mathfrak{D}_\mathfrak{n} + (n-2)!\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\big)\Big)_{\pi,\pi} = \Pi_n$, then $(X - \frac{n!}{2}\mathfrak{d}_\mathfrak{n})X$ resp. $X\big(X +$

$(n-2)!\mathfrak{d}_\mathfrak{n}\big)$ is not the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$.
We have

$$\Big(\big(\mathfrak{D}_\mathfrak{n} - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\big)\big(\mathfrak{D}_\mathfrak{n} + (n-2)!\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\big)\Big)_{\pi,\iota} = \Pi_n - \mathtt{des}_\mathtt{X}(\pi)\frac{n!}{2}\mathfrak{d}_\mathfrak{n}.$$

Since there are $\pi, \tau \in \mathcal{S}_n$ such that $\mathtt{des}_\mathtt{X}(\pi) \neq \mathtt{des}_\mathtt{X}(\tau)$, then

$$\Big(\big(\mathfrak{D}_\mathfrak{n} - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\big)\big(\mathfrak{D}_\mathfrak{n} + (n-2)!\mathfrak{d}_\mathfrak{n}\mathrm{I}_{n!}\big)\Big)_{\pi,\tau\in\mathcal{S}_n} \neq (0)_{\pi,\tau\in\mathcal{S}_n},$$

and $\big(X - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\big)\big(X + (n-2)!\mathfrak{d}_\mathfrak{n}\big)$ is not the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$.
We conclude that the minimal polynomial of $\mathfrak{D}_\mathfrak{n}$ is $X\big(X - \frac{n!}{2}\mathfrak{d}_\mathfrak{n}\big)\big(X + (n-2)!\mathfrak{d}_\mathfrak{n}\big)$. ☐

4 Minimal Polynomial of the Multinomial Inversion Statistic

To determine the minimal polynomial of $\mathfrak{I}_\mathfrak{n}$ we need to prove several multinomial formulas:
Let $n \geq 4$ and $i_1, j_1, i_2, j_2, a, b \in [n]$. We write:

$$\mathcal{S}_n^{i_1\,j_1} := \{\sigma \in \mathcal{S}_n \mid \sigma(i_1) > \sigma(j_1)\},$$

$$\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=a,\bar{j_2}=b} := \{\sigma \in \mathcal{S}_n^{i_1\,j_1} \mid \sigma^{-1}(i_2) = a,\ \sigma^{-1}(j_2) = b\}.$$

Lemma 4.1. *Let $n \geq 4$ and $i_1, j_1, i_2, j_2, a, b \in [n]$, with $i_1 < j_1$, $i_2 < j_2$. We have:*

- *if $\{a, b\} = \{i_1, j_1\}$:*
$$\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=j_1,\bar{j_2}=i_1} = \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=i_1,\bar{j_2}=j_1} = (n-2)!.$$
- *if $a, b \notin \{i_1, j_1\}$:*
$$\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=a,\bar{j_2}=b} = \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=a,\bar{j_2}=b} = \frac{(n-2)!}{2}.$$
- *if $a = i_1$ and $b \notin \{i_1, j_1\}$:*
$$\begin{aligned}\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=i_1,\bar{j_2}=b} &= (i_2-1)(n-3)!,\\ \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=i_1,\bar{j_2}=b} &= (n-i_2-1)(n-3)!.\end{aligned}$$
- *if $a \notin \{i_1, j_1\}$ and $b = i_1$:*
$$\begin{aligned}\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=a,\bar{j_2}=i_1} &= (j_2-2)(n-3)!,\\ \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=a,\bar{j_2}=i_1} &= (n-j_2)(n-3)!.\end{aligned}$$
- *if $a = j_1$ and $b \notin \{i_1, j_1\}$:*
$$\begin{aligned}\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=j_1,\bar{j_2}=b} &= (n-i_2-1)(n-3)!,\\ \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=j_1,\bar{j_2}=b} &= (i_2-1)(n-3)!.\end{aligned}$$
- *if $a \notin \{i_1, j_1\}$ and $b = j_1$:*
$$\begin{aligned}\#\mathcal{S}_n^{i_1\,j_1}{}_{|\bar{i_2}=a,\bar{j_2}=j_1} &= (n-j_2)(n-3)!,\\ \#\mathcal{S}_n^{j_1\,i_1}{}_{|\bar{i_2}=a,\bar{j_2}=j_1} &= (j_2-2)(n-3)!.\end{aligned}$$

Proof. Cardinality calculating. ☐

Let $n \geq 4$ and $i, j \in [n]$. We write

$${}^{i\,j}\mathcal{S}_n := \{\sigma \in \mathcal{S}_n \mid \sigma^{-1}(i) > \sigma^{-1}(j)\},$$

$$\chi_{i,j} = j - i - 1.$$

Lemma 4.2. *Let $n \geq 4$ and $i_1, j_1, i_2, j_2 \in [n]$ with $i_1 < j_1$ and $i_2 < j_2$. Then:*

(a)

$$\#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_2\,j_2} = (n-3)!\chi_{i_1,j_1}\chi_{i_2,j_2} + (n-2)!\frac{n^2-n+2}{4},$$

(b)

$$\#^{i_i\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{j_2\,i_2} = (n-2)!\frac{n^2-n-2}{4} - (n-3)!\chi_{i_1,j_1}\chi_{i_2,j_2}.$$

Proof. (a)

$$\begin{array}{rcc}
\#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_2\,j_2} & = & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) = j_1, \sigma(j_2) = i_1\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) = j_1, \sigma(j_2) \neq i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) > \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) \neq j_1, \sigma(j_2) = i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) > \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(j_2) = j_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) > \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) = i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) > \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) \notin \{i_1, j_1\}, \sigma(j_2) \notin \{i_1, j_1\}, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) > \sigma(j_2)\} \\
& = & (n-2)! \\
& + & (j_1-2)(n-i_2-1)(n-3)! \\
& + & (n-i_1-1)(j_2-2)(n-3)! \\
& + & (n-j_1)(n-j_2)(n-3)! \\
& + & (i_1-1)(i_2-1)(n-3)! \\
& + & \binom{n-2}{2}\binom{n-2}{2}(n-4)! \\
& = & (n-3)!\Big((i_1-j_1+1)(i_2-j_2+1) + \frac{n^3-3n^2+4n-4}{4}\Big).
\end{array}$$

(b)

$$\begin{array}{rcc}
\#^{i_i\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{j_2\,i_2} & = & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) = i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) < \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(j_2) = j_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) < \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) \neq j_1, \sigma(j_2) = i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) < \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) = j_1, \sigma(j_2) \neq i_1, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) < \sigma(j_2)\} \\
& + & \#\{\sigma \in \mathcal{S}_n \mid \sigma(i_2) \notin \{i_1, j_1\}, \sigma(j_2) \notin \{i_1, j_1\}, \sigma^{-1}(j_1) < \sigma^{-1}(i_1), \sigma(i_2) < \sigma(j_2)\} \\
& = & (n-i_1-1)(i_2-1)(n-3)! \\
& + & (j_1-2)(n-j_2)(n-3)! \\
& + & (i_1-1)(j_2-2)(n-3)! \\
& + & (n-j_1)(n-i_2-1)(n-3)! \\
& + & \binom{n-2}{2}\binom{n-2}{2}(n-4)! \\
& = & (n-3)!\Big((i_1-j_1+1)(-i_2+j_2-1) + \frac{(n+1)(n-2)^2}{4}\Big).
\end{array}$$

□

Let $n \geq 4$. We define:

$$\begin{array}{rccc}
\mathtt{f}_2: & \mathcal{S}_n & \to & \mathbb{R}[X_{1,2}, \ldots, X_{n-1,n}] \\
& \pi & \mapsto & \sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma^{-1})\mathtt{inv}_{\mathtt{X}}(\sigma\pi)
\end{array}$$

Lemma 4.3. *Let $n \geq 4$. Then:*

(a) For $(i_1, j_1) = (i_2, j_2)$:

▷ *If $\pi(i_1) < \pi(j_1)$:*

$$[X_{i_1,j_1}^2]\mathtt{f}_2(\pi) = (n-3)!\chi_{i_1,j_1}\chi_{\pi(i_1),\pi(j_1)} + (n-2)!\frac{n^2-n+2}{4}.$$

▷ *If $\pi(i_1) > \pi(j_1)$:*

$$[X_{i_1,j_1}^2]\mathtt{f}_2(\pi) = (n-2)!\frac{n^2-n-2}{4} - (n-3)!\chi_{i_1,j_1}\chi_{\pi(j_1),\pi(i_1)}.$$

(b) For $(i_1, j_1) \neq (i_2, j_2)$:

▷ *If* $\pi(i_1) < \pi(j_1)$ *and* $\pi(i_2) < \pi(j_2)$*:*

$$[X_{i_1,j_1}X_{i_2,j_2}]\mathtt{f}_2(\pi) = (n-3)!\big(\chi_{i_1,j_1}\chi_{\pi(i_2),\pi(j_2)} + \chi_{\pi(i_1),\pi(j_1)}\chi_{i_2,j_2}\big) + (n-2)!\frac{n^2-n+2}{2}.$$

▷ *If* $\pi(i_1) < \pi(j_1)$ *and* $\pi(i_2) > \pi(j_2)$*:*

$$[X_{i_1,j_1}X_{i_2,j_2}]\mathtt{f}_2(\pi) = (n-3)!\big(\chi_{\pi(i_1),\pi(j_1)}\chi_{i_2,j_2} - \chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)}\big) + \frac{n!}{2}.$$

▷ *If* $\pi(i_1) > \pi(j_1)$ *and* $\pi(i_2) > \pi(j_2)$*:*

$$[X_{i_1,j_1}X_{i_2,j_2}]\mathtt{f}_2(\pi) = (n-2)!\frac{n^2-n-2}{2} - (n-3)!\big(\chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)} + \chi_{\pi(j_1),\pi(i_1)}\chi_{i_2,j_2}\big).$$

Proof. (a) For $(i_1,j_1) = (i_2,j_2)$:

$$[X^2_{i_1,j_1}]\mathtt{f}_2(\pi) = \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}.$$

(b) For $(i_1,j_1) \neq (i_2,j_2)$:

$$[X_{i_1,j_1}X_{i_2,j_2}]\mathtt{f}_2(\pi) = \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{\pi(i_2)\,\pi(j_2)} + \#^{i_2\,j_2}\mathcal{S}_n \cap \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}.$$

□

Lemma 4.4. *Let* $n \geq 4$ *and* $i_1, j_1, i_2, j_2, i_3, j_3 \in [n]$ *with* $i_1 < j_1$*,* $i_2 < j_2$ *and* $i_3 < j_3$*. Then:*

(a)

$$\sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=b} = \frac{n!^2}{8} - \frac{(n-2)!^2}{2} - (n-4)(n-3)!^2\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{i_3,j_3}$$
$$-(n-3)!(n-2)!\Big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{i_3,j_3}\Big).$$

(b)

$$\sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i}_2=a,\bar{j}_2=b} = \frac{n!^2}{8} + \frac{(n-2)!^2}{2} + (n-4)(n-3)!^2\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{i_3,j_3}$$
$$+(n-3)!(n-2)!\Big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{i_3,j_3}\Big).$$

Proof. Using Lemma 4.1 and Lemma 4.2, we get:

(a)

$$\begin{aligned}
\textstyle\sum_{a,b\in[n]} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=b} &= \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{j_3\,i_3} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=j_3,\bar{j}_2=i_3} \\
&+ \#\textstyle\bigcup_{a,b\notin\{i_3,j_3\}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=b} \\
&+ \#\textstyle\bigcup_{b<i_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=i_3,\bar{j}_2=b} \\
&+ \#\textstyle\bigcup_{\substack{b>i_3\\ b\neq j_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=i_3,\bar{j}_2=b} \\
&+ \#\textstyle\bigcup_{\substack{b<j_3\\ b\neq i_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=i_3,\bar{j}_2=b} \\
&+ \#\textstyle\bigcup_{b>j_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{j_3\,b} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=j_3,\bar{j}_2=b} \\
&+ \#\textstyle\bigcup_{a<i_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,i_3} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=i_3} \\
&+ \#\textstyle\bigcup_{\substack{a>i_3\\ a\neq j_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,i_3} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=i_3} \\
&+ \#\textstyle\bigcup_{\substack{a<j_3\\ a\neq i_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,j_3} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=j_3} \\
&+ \#\textstyle\bigcup_{a>j_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,j_3} \times \mathcal{S}_n^{i_3\,j_3}{}_{|\bar{i}_2=a,\bar{j}_2=j_3}
\end{aligned}$$

$$
\begin{array}{cc}
= & (n-2)!^2\,\frac{n^2-n-2}{4} - (n-3)!(n-2)!\chi_{i_1,j_1}\chi_{i_3,j_3} \\
+ & \binom{n-2}{2}\frac{(n-2)!n!}{4} \\
+ & (i_2-1)(i_3-1)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (i_2-1)\binom{i_3-1}{2}(n-3)!^2\chi_{i_1,j_1} \\
+ & (i_2-1)\binom{n-i_3}{2}(n-3)!^2\chi_{i_1,j_1} - (i_2-1)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
 & +(i_2-1)(n-i_3-1)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (n-i_2-1)(j_3-2)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (n-i_2-1)\binom{j_3-1}{2}(n-3)!^2\chi_{i_1,j_1} \\
 & +(n-i_2-1)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
+ & (n-i_2-1)\binom{n-j_3}{2}(n-3)!^2\chi_{i_1,j_1} + (n-i_2-1)(n-j_3)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (j_2-2)\binom{i_3-1}{2}(n-3)!^2\chi_{i_1,j_1} + (j_2-2)(i_3-1)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (j_2-2)(n-i_3-1)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (j_2-2)\binom{n-i_3}{2}(n-3)!^2\chi_{i_1,j_1} \\
 & +(j_2-2)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
+ & (n-j_2)\binom{j_3-1}{2}(n-3)!^2\chi_{i_1,j_1} - (n-j_2)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
 & +(n-j_2)(j_3-2)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (n-j_2)(n-j_3)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (n-j_2)\binom{n-j_3}{2}(n-3)!^2\chi_{i_1,j_1} \\
= & \frac{n!^2}{8} - \frac{(n-2)!^2}{2} - (n-4)(n-3)!^2\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{i_3,j_3} \\
 & -(n-3)!(n-2)!\Big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{i_3,j_3}\Big).
\end{array}
$$

(b)

$$
\begin{array}{rcl}
\sum_{a,b\in[n]} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=b} & = & \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,j_3} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=i_3,\bar{j_2}=j_3} \\
 & + & \#\bigcup_{a,b\notin\{i_3,j_3\}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=b} \\
 & + & \#\bigcup_{b<i_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=i_3,\bar{j_2}=b} \\
 & + & \#\bigcup_{\substack{b>i_3\\ b\neq j_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=i_3,\bar{j_2}=b} \\
 & + & \#\bigcup_{\substack{b<j_3\\ b\neq i_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{i_3\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=i_3,\bar{j_2}=b} \\
 & + & \#\bigcup_{b>j_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{j_3\,b} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=j_3,\bar{j_2}=b} \\
 & + & \#\bigcup_{a<i_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,i_3} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=i_3} \\
 & + & \#\bigcup_{\substack{a>i_3\\ a\neq j_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,i_3} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=i_3} \\
 & + & \#\bigcup_{\substack{a<j_3\\ a\neq i_3}} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,j_3} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=j_3} \\
 & + & \#\bigcup_{a>j_3} {}^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,j_3} \times \mathcal{S}_n^{j_3\,i_3}{}_{|\bar{i_2}=a,\bar{j_2}=j_3}
\end{array}
$$

$$
\begin{array}{cc}
= & (n-3)!(n-2)!\chi_{i_1,j_1}\chi_{i_3,j_3} + (n-2)!^2\,\frac{n^2-n+2}{4} \\
+ & \binom{n-2}{2}\frac{(n-2)!n!}{4} \\
+ & (n-i_2-1)(i_3-1)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (n-i_2-1)\binom{i_3-1}{2}(n-3)!^2\chi_{i_1,j_1} \\
+ & (n-i_2-1)\binom{n-i_3}{2}(n-3)!^2\chi_{i_1,j_1} - (n-i_2-1)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
 & +(n-i_2-1)(n-i_3-1)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (i_2-1)(j_3-2)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (i_2-1)\binom{j_3-1}{2}(n-3)!^2\chi_{i_1,j_1} + (i_2-1)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
+ & (i_2-1)\binom{n-j_3}{2}(n-3)!^2\chi_{i_1,j_1} + (i_2-1)(n-j_3)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (n-j_2)\binom{i_3-1}{2}(n-3)!^2\chi_{i_1,j_1} + (n-j_2)(i_3-1)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (n-j_2)(n-i_3-1)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (n-j_2)\binom{n-i_3}{2}(n-3)!^2\chi_{i_1,j_1} \\
 & +(n-j_2)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} \\
+ & (j_2-2)\binom{j_3-1}{2}(n-3)!^2\chi_{i_1,j_1} - (j_2-2)(n-3)!^2\chi_{i_1,j_1}\chi_{i_3,j_3} + (j_2-2)(j_3-2)(n-3)!(n-2)!\frac{n^2-n+2}{4} \\
+ & (j_2-2)(n-j_3)(n-3)!(n-2)!\frac{n^2-n-2}{4} - (j_2-2)\binom{n-j_3}{2}(n-3)!^2\chi_{i_1,j_1} \\
= & \frac{n!^2}{8} + \frac{(n-2)!^2}{2} + (n-4)(n-3)!^2\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{i_3,j_3} \\
 & +(n-3)!(n-2)!\Big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{i_3,j_3}\Big)
\end{array}
$$

□

Let $n \geq 4$. We define:

$$
\begin{array}{rccc}
\mathtt{f_3}: & \mathcal{S}_n & \to & \mathbb{R}[X_{1,2},\ldots,X_{n-1,n}] \\
 & \pi & \mapsto & \sum_{\sigma,\tau\in\mathcal{S}_n} \mathtt{inv_X}(\sigma^{-1})\mathtt{inv_X}(\sigma\tau^{-1})\mathtt{inv_X}(\tau\pi).
\end{array}
$$

Lemma 4.5. *Let $n \geq 4$. Then:*

(a) For $(i_1,j_1) = (i_2,j_2) = (i_3,j_3)$:

▷ *If $\pi(i_1) < \pi(j_1)$:*

$$[X^3_{i_1,j_1}]\mathtt{f3}(\pi) = \frac{n!^2}{8} - \frac{(n-2)!^2}{2} - (n-4)(n-3)!^2\chi^2_{i_1,j_1}\chi_{\pi(i_1),\pi(j_1)}$$

$$-(n-3)!(n-2)!\chi^2_{i_1,j_1} - 2(n-3)!(n-2)!\chi_{i_1,j_1}\chi_{\pi(i_1),\pi(j_1)}.$$

▷ *If $\pi(i_1) > \pi(j_1)$:*

$$[X^3_{i_1,j_1}]\mathtt{f3}(\pi) = \frac{n!^2}{8} + \frac{(n-2)!^2}{2} + (n-4)(n-3)!^2\chi^2_{i_1,j_1}\chi_{\pi(j_1),\pi(i_1)}$$

$$+(n-3)!(n-2)!\chi^2_{i_1,j_1} + 2(n-3)!(n-2)!\chi_{i_1,j_1}\chi_{\pi(j_1),\pi(i_1)}.$$

(b) For $(i_1,j_1) = (i_2,j_2) \neq (i_3,j_3)$:

▷ *If $\pi(i_1) < \pi(j_1)$ and $\pi(i_3) < \pi(j_3)$:*

$$[X^2_{i_1,j_1}X_{i_3,j_3}]\mathtt{f3}(\pi) = \frac{3n!^2}{8} - \frac{3(n-2)!^2}{2} - (n-4)(n-3)!^2\big(\chi^2_{i_1,j_1}\chi_{\pi(i_3),\pi(j_3)} + 2\chi_{i_1,j_1}\chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big)$$

$$-(n-3)!(n-2)!\big(\chi^2_{i_1,j_1} + 2\chi_{i_1,j_1}\chi_{\pi(i_3),\pi(j_3)} + 2\chi_{i_1,j_1}\chi_{i_3,j_3} + 2\chi_{i_1,j_1}\chi_{\pi(i_1),\pi(j_1)} + 2\chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big).$$

▷ *If $\pi(i_1) < \pi(j_1)$ and $\pi(i_3) > \pi(j_3)$:*

$$[X^2_{i_1,j_1}X_{i_3,j_3}]\mathtt{f3}(\pi) = \frac{3n!^2}{8} - \frac{(n-2)!^2}{2} + (n-4)(n-3)!^2\big(\chi^2_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} - 2\chi_{i_1,j_1}\chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big)$$

$$+(n-3)!(n-2)!\big(\chi^2_{i_1,j_1} + 2\chi_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} - 2\chi_{i_1,j_1}\chi_{i_3,j_3} - 2\chi_{i_1,j_1}\chi_{\pi(i_1),\pi(j_1)} - 2\chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big).$$

▷ *If $\pi(i_1) > \pi(j_1)$ and $\pi(i_3) < \pi(j_3)$:*

$$[X^2_{i_1,j_1}X_{i_3,j_3}]\mathtt{f3}(\pi) = \frac{3n!^2}{8} + \frac{(n-2)!^2}{2} - (n-4)(n-3)!^2\big(\chi^2_{i_1,j_1}\chi_{\pi(i_3),\pi(j_3)} - 2\chi_{i_1,j_1}\chi_{i_3,j_3}\chi_{\pi(j_1),\pi(i_1)}\big)$$

$$-(n-3)!(n-2)!\big(\chi^2_{i_1,j_1} + 2\chi_{i_1,j_1}\chi_{\pi(i_3),\pi(j_3)} - 2\chi_{i_1,j_1}\chi_{i_3,j_3} - 2\chi_{i_1,j_1}\chi_{\pi(j_1),\pi(i_1)} - 2\chi_{i_3,j_3}\chi_{\pi(j_1),\pi(i_1)}\big).$$

▷ *If $\pi(i_1) > \pi(j_1)$ and $\pi(i_3) > \pi(j_3)$:*

$$[X^2_{i_1,j_1}X_{i_3,j_3}\mathtt{f3}(\pi) = \frac{3n!^2}{8} + \frac{3(n-2)!^2}{2} + (n-4)(n-3)!^2\big(\chi^2_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} + 2\chi_{i_1,j_1}\chi_{i_3,j_3}\chi_{\pi(j_1),\pi(i_1)}\big)$$

$$+(n-3)!(n-2)!\big(\chi^2_{i_1,j_1} + 2\chi_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} + 2\chi_{i_1,j_1}\chi_{i_3,j_3} + 2\chi_{i_1,j_1}\chi_{\pi(j_1),\pi(i_1)} + 2\chi_{i_3,j_3}\chi_{\pi(j_1),\pi(i_1)}\big).$$

(c) For $(i_1,j_1) \neq (i_2,j_2) \neq (i_3,j_3)$:

▷ *If $\pi(i_1) < \pi(j_1)$, $\pi(i_2) < \pi(j_2)$ and $\pi(i_3) < \pi(j_3)$:*

$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f3}(\pi) = \frac{3n!^2}{4} - 3(n-2)!^2$$

$$-2(n-4)(n-3)!^2\big(\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{\pi(i_3),\pi(j_3)} + \chi_{i_1,j_1}\chi_{\pi(i_2),\pi(j_2)}\chi_{i_3,j_3} + \chi_{\pi(i_1),\pi(j_1)}\chi_{i_2,j_2}\chi_{i_3,j_3}\big)$$

$$-2(n-3)!(n-2)!\big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{\pi(i_3),\pi(j_3)} + \chi_{i_2,j_2}\chi_{\pi(i_3),\pi(j_3)}$$

$$+\chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_1,j_1}\chi_{\pi(i_2),\pi(j_2)} + \chi_{i_3,j_3}\chi_{\pi(i_2),\pi(j_2)} + \chi_{i_2,j_2}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{\pi(i_1),\pi(j_1)} + \chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big).$$

▷ *If $\pi(i_1) < \pi(j_1)$, $\pi(i_2) < \pi(j_2)$ and $\pi(i_3) > \pi(j_3)$:*

$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f3}(\pi) = \frac{3n!^2}{4} - (n-2)!^2$$

$$+2(n-4)(n-3)!^2\big(\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)} - \chi_{i_1,j_1}\chi_{\pi(i_2),\pi(j_2)}\chi_{i_3,j_3} - \chi_{\pi(i_1),\pi(j_1)}\chi_{i_2,j_2}\chi_{i_3,j_3}\big)$$

$$+2(n-3)!(n-2)!\big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} + \chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)}$$

$$-\chi_{i_1,j_1}\chi_{i_3,j_3} - \chi_{i_1,j_1}\chi_{\pi(i_2),\pi(j_2)} - \chi_{i_3,j_3}\chi_{\pi(i_2),\pi(j_2)} - \chi_{i_2,j_2}\chi_{i_3,j_3} - \chi_{i_2,j_2}\chi_{\pi(i_1),\pi(j_1)} - \chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big).$$

▷ *If $\pi(i_1) < \pi(j_1)$, $\pi(i_2) > \pi(j_2)$ and $\pi(i_3) > \pi(j_3)$:*

$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f_3}(\pi) = \frac{3n!^2}{4} + (n-2)!^2$$
$$+2(n-4)(n-3)!^2\big(\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)} + \chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)}\chi_{i_3,j_3} - \chi_{\pi(i_1),\pi(j_1)}\chi_{i_2,j_2}\chi_{i_3,j_3}\big)$$
$$+2(n-3)!(n-2)!\big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} + \chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)}$$
$$+\chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)} + \chi_{i_3,j_3}\chi_{\pi(j_2),\pi(i_2)} - \chi_{i_2,j_2}\chi_{i_3,j_3} - \chi_{i_2,j_2}\chi_{\pi(i_1),\pi(j_1)} - \chi_{i_3,j_3}\chi_{\pi(i_1),\pi(j_1)}\big).$$

▷ *If $\pi(i_1) > \pi(j_1)$, $\pi(i_2) > \pi(j_2)$ and $\pi(i_3) > \pi(j_3)$:*

$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f_3}(\pi) = \frac{3n!^2}{4} + 3(n-2)!^2$$
$$+2(n-4)(n-3)!^2\big(\chi_{i_1,j_1}\chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)} + \chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)}\chi_{i_3,j_3} + \chi_{\pi(j_1),\pi(i_1)}\chi_{i_2,j_2}\chi_{i_3,j_3}\big)$$
$$+2(n-3)!(n-2)!\big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{\pi(j_3),\pi(i_3)} + \chi_{i_2,j_2}\chi_{\pi(j_3),\pi(i_3)}$$
$$+\chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_1,j_1}\chi_{\pi(j_2),\pi(i_2)} + \chi_{i_3,j_3}\chi_{\pi(j_2),\pi(i_2)} + \chi_{i_2,j_2}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{\pi(j_1),\pi(i_1)} + \chi_{i_3,j_3}\chi_{\pi(j_1),\pi(i_1)}\big).$$

Proof. (a) For $(i_1,j_1) = (i_2,j_2) = (i_3,j_3)$:

$$[X_{i_1,j_1}^3]\mathtt{f_3}(\pi) = \sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}{}_{|\bar{i_1}=a,\bar{j_1}=b}.$$

(b) For $(i_1,j_1) = (i_2,j_2) \neq (i_3,j_3)$:

$$[X_{i_1,j_1}^2 X_{i_3,j_3}]\mathtt{f_3}(\pi) = \sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_3)\,\pi(j_3)}{}_{|\bar{i_1}=a,\bar{j_1}=b}$$
$$+ \sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}{}_{|\bar{i_3}=a,\bar{j_3}=b} + \sum_{a,b\in[n]} \#^{i_3\,j_3}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}{}_{|\bar{i_1}=a,\bar{j_1}=b}.$$

(c) For $(i_1,j_1) \neq (i_3,j_3) \neq (i_2,j_2)$:

$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f_3}(\pi) = \sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_3)\,\pi(j_3)}{}_{|\bar{i_2}=a,\bar{j_2}=b}$$
$$+ \sum_{a,b\in[n]} \#^{i_2\,j_2}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_3)\,\pi(j_3)}{}_{|\bar{i_1}=a,\bar{j_1}=b} + \sum_{a,b\in[n]} \#^{i_1\,j_1}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_2)\,\pi(j_2)}{}_{|\bar{i_3}=a,\bar{j_3}=b}$$
$$+ \sum_{a,b\in[n]} \#^{i_3\,j_3}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_2)\,\pi(j_2)}{}_{|\bar{i_1}=a,\bar{j_1}=b} + \sum_{a,b\in[n]} \#^{i_2\,j_2}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}{}_{|\bar{i_3}=a,\bar{j_3}=b}$$
$$+ \sum_{a,b\in[n]} \#^{i_3\,j_3}\mathcal{S}_n \cap \mathcal{S}_n^{a\,b} \times \mathcal{S}_n^{\pi(i_1)\,\pi(j_1)}{}_{|\bar{i_2}=a,\bar{j_2}=b}.$$

□

Let $n \geq 4$. We write

$$\Lambda = (n-2)! \sum_{\{(i,j)\in[n]^2 \;|\; i<j\}} (j-i)X_{i,j},$$
$$\Delta = (n-3)! \sum_{\{(i,j)\in[n]^2 \;|\; i<j\}} \Big(n - 2(j-i)\Big)X_{i,j}.$$

Lemma 4.6. *Let $n \geq 4$. Then:*

(a)

$$[X_{i,j}](\Lambda + \Delta) = 2(n-2)! + (n-4)(n-3)!\chi_{i,j}.$$

(b)

$$[X_{i_1,j_1}X_{i_2,j_2}](\Lambda\Delta) = (n-3)!(n-2)!\frac{\big(2(n-2) + (n-4)\chi_{i_1,j_1} + (n-4)\chi_{i_2,j_2} - 4\chi_{i_1,j_1}\chi_{i_2,j_2}\big)}{1 + \delta_{(i_1,j_1),(i_2,j_2)}}$$

Proof. Coefficient calculating. □

Let $n \geq 4$. We define:

$$\begin{array}{rccc} \mathtt{f}: & \mathcal{S}_n & \rightarrow & \mathbb{R}[X_{1,2}, \ldots, X_{n-1,n}] \\ & \pi & \mapsto & \Lambda\Delta\mathtt{inv}_{\mathtt{X}}(\pi) + \big(\Lambda + \Delta\big)\mathtt{f}_2(\pi) + \mathtt{f}_3(\pi) \end{array}$$

We write ι for the identity permutation of $\mathcal{S}_n$.

Lemma 4.7. *Let $n \geq 4$ and $\pi \in \mathcal{S}_n$. Then:*

$$\mathtt{f}(\pi) = \mathtt{f}(\iota) = \big(\Lambda + \Delta\big)\mathtt{f}_2(\iota) + \mathtt{f}_3(\iota).$$

Proof. Using Lemma 4.3, Lemma 4.5 and Lemma 4.6, we prove that:

- for $(i_1, j_1) = (i_2, j_2) = (i_3, j_3)$:
$$[X_{i_1,j_1}^3]\mathtt{f}(\iota) = [X_{i_1,j_1}^3]\mathtt{f}(\pi)$$
$$= \frac{n!^2}{8} + (n-2)!^2\frac{n^2-n+1}{2} + (n-4)(n-3)!(n-2)!\frac{n^2-n+2}{4}\chi_{i_1,j_1} - (n-3)!(n-2)!\chi_{i_1,j_1}^2,$$
- for $(i_1, j_1) = (i_2, j_2) \neq (i_3, j_3)$:
$$[X_{i_1,j_1}^2 X_{i_3,j_3}]\mathtt{f}(\iota) = [X_{i_1,j_1}^2 X_{i_3,j_3}]\mathtt{f}(\pi)$$
$$= \frac{3n!^2}{8} + \frac{3(n-2)!n!}{2} + \frac{3(n-2)!^2}{2}$$
$$+(n-4)(n-3)!(n-2)!\frac{n^2-n+2}{4}\big(2\chi_{i_1,j_1} + \chi_{i_3,j_3}\big)$$
$$-(n-3)!(n-2)!\big(\chi_{i_1,j_1}^2 + 2\chi_{i_1,j_1}\chi_{i_3,j_3}\big),$$
- for $(i_1, j_1) \neq (i_2, j_2) \neq (i_3, j_3) \neq (i_1, j_1)$:
$$[X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f}(\iota) = [X_{i_1,j_1}X_{i_2,j_2}X_{i_3,j_3}]\mathtt{f}(\pi)$$
$$= \frac{3n!^2}{4} + 3(n-2)!n! + 3(n-2)!^2$$
$$+(n-4)(n-3)!(n-2)!\frac{n^2-n+2}{2}\big(\chi_{i_1,j_1} + \chi_{i_2,j_2} + \chi_{i_3,j_3}\big)$$
$$-2(n-3)!(n-2)!\big(\chi_{i_1,j_1}\chi_{i_2,j_2} + \chi_{i_1,j_1}\chi_{i_3,j_3} + \chi_{i_2,j_2}\chi_{i_3,j_3}\big).$$

□

We are now in position to determine the minimal polynomial of $\mathfrak{I}_{\mathtt{n}}$.
Let $n \geq 4$. We write

$$\Omega = \frac{n!}{2} \sum_{\{(i,j)\in[n]^2 \ | \ i<j\}} X_{i,j}.$$

Proposition 4.8. *Let $n \geq 4$. The minimal polynomial of $\mathfrak{I}_{\mathtt{n}}$ is:*

$$X\big(X + \Lambda\big)\big(X + \Delta\big)\big(X - \Omega\big).$$

Proof. Using Lemma 2.4 and Lemma 4.7, we get:

$$\big(\sum_{\sigma\in\mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Lambda\iota\big)\big(\sum_{\sigma\in\mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Delta\iota\big)\sum_{\sigma\in\mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma$$

$$= \sum_{\sigma\in\mathcal{S}_n}\big(\sum_{\substack{\sigma_1,\sigma_2,\sigma_3\in\mathcal{S}_n \\ \sigma_1\sigma_2\sigma_3=\sigma}} \mathtt{inv}_{\mathtt{X}}(\sigma_1)\mathtt{inv}_{\mathtt{X}}(\sigma_2)\mathtt{inv}_{\mathtt{X}}(\sigma_3)\big)\sigma + \sum_{\sigma\in\mathcal{S}_n}\big((\Lambda+\Delta)\sum_{\substack{\sigma_1,\sigma_2\in\mathcal{S}_n \\ \sigma_1\sigma_2=\sigma}} \mathtt{inv}_{\mathtt{X}}(\sigma_1)\mathtt{inv}_{\mathtt{X}}(\sigma_2)\big)\sigma + \sum_{\sigma\in\mathcal{S}_n\setminus\{\iota\}} \Lambda\Delta\mathtt{inv}_{\mathtt{X}}(\sigma)\sigma$$

$$= \sum_{\sigma\in\mathcal{S}_n}\big(\sum_{\sigma_1,\sigma_2\in\mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma_1^{-1})\mathtt{inv}_{\mathtt{X}}(\sigma_1\sigma_2^{-1})\mathtt{inv}_{\mathtt{X}}(\sigma_2\sigma)\big)\sigma + \sum_{\sigma\in\mathcal{S}_n}\big((\Lambda+\Delta)\sum_{\sigma_1\in\mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma_1^{-1})\mathtt{inv}_{\mathtt{X}}(\sigma_1\sigma)\big)\sigma + \sum_{\sigma\in\mathcal{S}_n\setminus\{\iota\}} \Lambda\Delta\mathtt{inv}_{\mathtt{X}}(\sigma)\sigma$$

$$= \sum_{\sigma \in \mathcal{S}_n} \mathtt{f}(\sigma)\sigma = \mathtt{f}(\iota) \sum_{\sigma \in \mathcal{S}_n} \sigma.$$

Then:

$$\Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Lambda\iota \Big) \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Delta\iota \Big) \sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma - \Omega\iota \Big)$$

$$= \mathtt{f}(\iota) \sum_{\sigma \in \mathcal{S}_n} \sigma \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma - \Omega\iota \Big) = 0.$$

Hence the minimal polynomial of $\mathfrak{I}_{\mathfrak{n}}$ divides $X\big(X+\Lambda\big)\big(X+\Delta\big)\big(X-\Omega\big)$.
It is clear that the minimal polynomial of $\mathfrak{I}_{\mathfrak{n}}$ does not divide $X\big(X+\Lambda\big)\big(X+\Delta\big)$.
We have:

$$[X_{1,4}^3]\Bigg([\iota]\Big(\Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Lambda\iota \Big) \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma - \Omega\iota \Big) \sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma \Big) \Bigg)$$

$$= (n-3)!^2 \frac{n^4 - 8n^3 + 22n^2 - 36n + 44}{2} \neq 0.$$

Then the minimal polynomial of $\mathfrak{I}_{\mathfrak{n}}$ does not divide $X\big(X+\Lambda\big)\big(X-\Omega\big)$.
We have:

$$[X_{1,3}^3]\Bigg([\iota]\Big(\Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Delta\iota \Big) \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma - \Omega\iota \Big) \sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma \Big) \Bigg)$$

$$= -4(n-3)!(n-2)! - (n-3)!n! \neq 0.$$

Then the minimal polynomial of $\mathfrak{I}_{\mathfrak{n}}$ does not divide $X\big(X+\Delta\big)\big(X-\Omega\big)$.
We have:

$$[X_{1,3}^3]\Bigg([\iota]\Big(\Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Lambda\iota \Big) \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma + \Delta\iota \Big) \Big(\sum_{\sigma \in \mathcal{S}_n} \mathtt{inv}_{\mathtt{X}}(\sigma)\sigma - \Omega\iota \Big) \Big) \Bigg)$$

$$= \frac{(n-2)!^2}{2} + \frac{(n-2)!n!}{2} - \frac{(n-2)!^2 n!}{2} \neq 0.$$

Then the minimal polynomial of $\mathfrak{I}_{\mathfrak{n}}$ does not divide $\big(X+\Lambda\big)\big(X+\Delta\big)\big(X-\Omega\big)$. □

5 Proofs of the Theorems

We prove first Theorem 1 and then Theorem 2.

Proof. From Proposition 3.3, we deduce that $\mathfrak{D}_{\mathfrak{n}}$ is diagonalizable and:

$$Sp(\mathfrak{D}_{\mathfrak{n}}) = \{ \frac{n!}{2}\mathfrak{d}_{\mathfrak{n}},\ -(n-2)!\mathfrak{d}_{\mathfrak{n}},\ 0 \}.$$

From the diagonalisability of $\mathfrak{D}_{\mathfrak{n}}$ and the result (1) of Section 2, we get

$$V_{\mathfrak{D}_{\mathfrak{n}}}\big(\frac{n!}{2}\mathfrak{d}_{\mathfrak{n}}\big) = 1$$

.

The trace of $\mathfrak{D}_{\mathfrak{n}}$ is 0. Then:

$$\frac{n!}{2}\mathfrak{d}_{\mathfrak{n}}\, V_{\mathfrak{D}_{\mathfrak{n}}}\big(\frac{n!}{2}\mathfrak{d}_{\mathfrak{n}}\big) - (n-2)!\mathfrak{d}_{\mathfrak{n}}\, V_{\mathfrak{D}_{\mathfrak{n}}}\big(-(n-2)!\mathfrak{d}_{\mathfrak{n}}\big) + 0\, V_{\mathfrak{D}_{\mathfrak{n}}}(0) = 0$$

$$V_{\mathfrak{D}_{\mathfrak{n}}}\big(-(n-2)!\mathfrak{d}_{\mathfrak{n}}\big) = \binom{n}{2}.$$

The dimension of $\mathfrak{D}_{\mathfrak{n}}$ is $n!$. Then:

$$V_{\mathfrak{D}_{\mathfrak{n}}}\big(\frac{n!}{2}\mathfrak{d}_{\mathfrak{n}}\big) + V_{\mathfrak{D}_{\mathfrak{n}}}\big(-(n-2)!\mathfrak{d}_{\mathfrak{n}}\big) + V_{\mathfrak{D}_{\mathfrak{n}}}(0) = n!$$

$$V_{\mathfrak{D}_{\mathfrak{n}}}(0) = n! - \binom{n}{2} - 1.$$

□

Proof. From Proposition 4.8, we deduce that $\mathfrak{I}_\mathfrak{n}$ is diagonalizable and:

$$Sp(\mathfrak{I}_\mathfrak{n}) = \{\Omega, -\Lambda, -\Delta, 0\}.$$

From the diagonalisability of $\mathfrak{I}_\mathfrak{n}$ and the result (2) of Section 2, we get

$$V_{\mathfrak{I}_\mathfrak{n}}(\Omega) = 1$$

.

The trace of $\mathfrak{I}_\mathfrak{n}$ is 0. Then:

- $$V_{\mathfrak{I}_\mathfrak{n}}(\Omega)\big(\Omega\,,\, X_{1,2}\big) + V_{\mathfrak{I}_\mathfrak{n}}(\Lambda)\big(\Lambda\,,\, X_{1,2}\big) + V_{\mathfrak{I}_\mathfrak{n}}(\Delta)\big(\Delta\,,\, X_{1,2}\big) + V_{\mathfrak{I}_\mathfrak{n}}(0)\big(0\,,\, X_{1,2}\big) = 0,$$
  $$V_{\mathfrak{I}_\mathfrak{n}}(\Lambda) + V_{\mathfrak{I}_\mathfrak{n}}(\Delta) = \binom{n}{2}.$$
- $$V_{\mathfrak{I}_\mathfrak{n}}(\Omega)\big(\Omega\,,\, X_{1,n}\big) + V_{\mathfrak{I}_\mathfrak{n}}(\Lambda)\big(\Lambda\,,\, X_{1,n}\big) + V_{\mathfrak{I}_\mathfrak{n}}(\Delta)\big(\Delta\,,\, X_{1,n}\big) + V_{\mathfrak{I}_\mathfrak{n}}(0)\big(0\,,\, X_{1,n}\big) = 0,$$
  $$(n-1)V_{\mathfrak{I}_\mathfrak{n}}(\Lambda) - V_{\mathfrak{I}_\mathfrak{n}}(\Delta) = \binom{n}{2}.$$

So we deduce:

$$V_{\mathfrak{I}_\mathfrak{n}}(\Lambda) = n-1 \text{ and } V_{\mathfrak{I}_\mathfrak{n}}(\Delta) = \binom{n-1}{2}.$$

The dimension of $\mathfrak{I}_\mathfrak{n}$ is $n!$. Then:

$$V_{\mathfrak{I}_\mathfrak{n}}(\Omega) + V_{\mathfrak{I}_\mathfrak{n}}(\Lambda) + V_{\mathfrak{I}_\mathfrak{n}}(\Delta) + V_{\mathfrak{I}_\mathfrak{n}}(0) = n!,$$

$$V_{\mathfrak{I}_\mathfrak{n}}(0) = n! - \binom{n}{2} - 1.$$

□

References

[1] G. Denham, and Ph. Hanlon, *Some Algebraic Properties of the Schechtman-Varchenko Bilinear Form*, MSRI Publications **38** (1999) 149-176

[2] I. M. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. S. Retakh, and J.-Y. Thibon, *Noncommutative Symmetric Functions*, Advances in Mathematics **112** (1995) 218-348

[3] R. Horn, and C. Johnson, *Matrix Analysis*, Cambridge University Press (1985)

[4] C. Krattenthaler, *Advanced Determinant Calculus*, Séminaire Lotharingien de Combinatoire **42** (1999) Article B42q, 67 pp

[5] D. Krob, B. Leclerc, and J.-Y. Thibon, *Noncommutative Symmetric Functions II: Transformations of Alphabets*, International Journal of Algebra and Computation **7** (1997) 181-264

[6] V. Reiner, F. Saliola, and V. Welker, *Spectra of Symmetrized Shuffling operators*, http://arxiv.org/abs/1102.2460 (2011)

[7] P. Renteln, *The Distance Spectra of Cayley Graphs of Coxeter Groups*, Discrete Mathematics **311** (2011) 738-755

[8] C. Reutenauer, *Free Lie Algebras*, Oxford Science Publications (1993)

[9] R. Stanley, *Enumerative Combinatorics*, Cambridge University Press (1997)

[10] A. Varchenko, *Bilinear Form of Real Configuration of Hyperplanes*, Advances in Mathematics **97** (1993) 110-144

[11] D. Zagier, *Realizability of a Model in Infinite Statistics*, Communications in Mathematical Physics **147** (1992) 199-210